

Table 1: Patients' demographic characteristics and treatment related parameters in group A (Conventional Radiotherapy treatment) and Group B (Hypofractionated Radiotherapy treatment).

Characteristics	All Patients	Group A	Group B
Age			
Mean	55.16	57.06	53.2
Range	31-69	42-69	31-67
Menopausal status			
Pre	18 (29%)	7 (11.5%)	11 (18%)
Post	43 (70.5%)	24 (39.3%)	19 (31.1%)
Left	38 (62.3%)	18 (29.5%)	20 (32.8%)
Right	23 (37.7%)	13 (21.3%)	10 (16.4%)
T Stage			
T1	48(78.7%)	25 (41%)	23 (37.7%)
T2	13 (21.3%)	6 (9.8%)	7 (11.5%)
Histology			
Ductal	60 (98.4%)	30 (49.2%)	30 (49.2%)
Lobular	1 (1.6%)	1 (3.2%)	0 (0%)
Grade			
1	38	20	18
2	20	10	10
3	3	1	2
ER Status			
+	56 (91.8%)	27 (44.3%)	29 (47.5%)
-	5 (8.2%)	4 (6.6%)	1 (1.6%)
PR Status			
+	48 (78.7%)	22 (36.1%)	26 (42.6%)
-	13 (21.3%)	9 (14.8%)	4 (6.6%)
HER2 Status			
+	10 (16.4%)	6 (9.8%)	4 (6.6%)
-	51 (83.6%)	25 (41%)	26 (42.6%)
Chemotherapy			
Yes	16(26.2%)	6 (9.8%)	10 (16.4%)
Hormonotherapy			
Yes			
AI	38 (62.3%)	20 (32.8%)	18 (29.5%)
Tamoxifene	18 (29.5%)	7(11.5%)	11 (18%)
No	5 (8.1%)	4(6.5%)	1 (1.6%)
Smoking			
Yes	25 (41%)	11 (18%)	14 (23%)
No	36 (59%)	20 (32.8%)	16 (26.2%)

AI: Aromatase Inhibitors