

Table 1: Antimicrobial susceptibility testing results of 83 isolates of *K. pneumoniae* collected from Mofid Children and Taleghani Hospitals, Tehran, Iran.

Antibiotic	Resistant No (%)	Intermediate No (%)	Sensitive No (%)
Aztreonam (10 µg)	49 (59%)	3 (3.6%)	31 (37.3%)
Meropenem (10 µg)	20 (24%)	2 (2.4%)	61 (73.5%)
Gentamicin (10 µg)	29 (35%)	3 (3.6%)	51 (61.5%)
Ciprofloxacin (30 µg)	46 (55.5%)	4 (4.8%)	33 (39.7%)
Amikacin (30 µg)	12 (14.4%)	4 (4.8%)	67 (80.7%)
Ceftazidime (30 µg)	46 (55.4%)	3 (3.6%)	34 (41%)
Imipenem (10 µg)	20 (24%)	2 (2.4%)	61 (73.5%)
Cefotaxime (30 µg)	50 (60.2)	2 (2.4%)	31 (37.3%)
Cefepime (FEP, 30 µg)	30 (36.15%)	9 (10.8%)	44 (53.05%)
Tetracycline (TE,10 µg)	33 (39.7%)	3 (3.6%)	57 (68.6%)
Ampicillin (AMP,10 µg)	65 (78.3%)	2 (2.4%)	16 (19.2%)
Piperacillin (PIP,100 µg)	50 (60.2%)	0 (0.0%)	33 (39.7%)
Ceftriaxone (CRO,30 µg)	49 (59%)	2 (2.4%)	32 (38.5%)
Cefpodoxime (CPD,30µg)	57 (68.6%)	3 (3.6%)	23 (27.7%)
Tigecycline (TGC,15µg)	5 (6.02%)	25 (30.1%)	53 (63.8%)
Doripenem (DOR,10 µg)	20 (24%)	1 (1.2%)	62 (74.6%)
Ertapenem (ETP, 10 µg)	21 (25.3%)	2 (2.4%)	57 (68.6%)
Piperacillin/Tazobactam (PTZ,100/10µg)	22 (26.5%)	5 (6.02%)	56 (67.4%)
Fosfomicin/Trometamol (FOT, 200 µg)	3 (3.6%)	12 (14.4%)	68 (82%)