

Table 5: Daily warfarin dose and clinical characteristics in patients with various *factor VII* genotypes.

	Wild Type (1) (GG)	Heterozygous (2) (GT)	Homozygous (3) (TT)	p Value	Multiple Comparison
	Mean \pm SD median (25th-75th)	Mean \pm SD median (25th-75th)	Mean \pm SD median (25th-75th)		
Age	60.76 \pm 13.11 64 (48-72)	65 \pm 10.30 65 (57.5-74.5)	63.76 \pm 11.57 64 (57-72)	0.450 ^a	-
Dose	5.09 \pm 1.69 5 (3.93-6.96)	4.06 \pm 1.74 3.93 (2.5-5)	3.74 \pm 1.20 3.57 (2.5-5)	0.038 ^b	1-3
BMI (kg/m ²)	24.19 \pm 4.58 23.33 (20.54-27.81)	25.25 \pm 2.99 25.53 (23.05-27.10)	25.82 \pm 3.79 24.90 (23.19-27.71)	0.501 ^a	-
INR	2.43 \pm 0.33 2.41 (2.13-2.77)	2.39 \pm 0.68 2.40 (2.01-2.79)	2.42 \pm 0.83 2.37 (1.86-2.66)	0.905 ^b	-
Sex^c	number (%)	number (%)	number (%)		
Female	27 (45%)	20 (33.3 %)	13 (21.7%)	0.277	
Male	20 (48.8%)	17 (41.5%)	4 (9.8%)		

SD: standard deviation, *factor VII*: coagulation *factor VII* genotype, *a*: one-way analysis of variance, *b*: Kruskal-Wallis Test, *c*: Chi-square test for sex, ** p \leq 0.001.