

**Table 4:** The one-trait-bi-method matrix with Spearman's correlation coefficients.

	MFS-T2	IPSA-T2	BIFS-T2	MAS-T2	MFS-T1	IPSA-T1	BIFS-T1	MAS-T1
MFS-T2	1							
IPSA-T2	0.391*	1						
BIFS-T2	0.502*	0.332*	1					
MAS-T2	0.396*	0.794*	0.339*	1				
MFS-T1	0.838*	0.320*	0.445*	0.326*	1			
IPSA-T1	0.357*	0.786*	0.281*	0.668*	0.343*	1		
BIFS-T1	0.673*	0.382*	0.583*	0.397*	0.622*	0.401*	1	
MAS-T1	0.310*	0.660*	0.267*	0.718*	0.294*	0.718*	0.357*	1

\*: significant correlation,  $p < 0.001$ . MFS-T2: Medical Fear Survey, filled in by the study participants themselves; IPSA-T2: Injection Phobia Scale-Anxiety, filled in by the study participants themselves; BIFS-T2: Blood/Injection Fear Scale, filled in by the study participants themselves; MAS-T2: Medical Avoidance Survey, filled in by the study participants themselves; MFS-T1: Medical Fear Survey, filled in by the study investigators; IPSA-T1: Injection Phobia Scale-Anxiety, filled in by the study investigators; BIFS-T1: Blood/Injection Fear Scale, filled in by the study investigators; MAS-T1: Medical Avoidance Survey, filled in by the study investigators.