

Table 1: Thyroid cancer incidence rates in Serbia, per 100,000 residents, for females and males during the period 1999-2008.

Age group	0-19	20-29	30-39	40-49	50-59	60-69	70+	All ages	
Years								CR	AAR
Females									
1999	0.30	0.28	1.94	4.11	5.64	4.72	4.20	2.72	1.89
2000	0.32	0.55	2.80	5.33	2.32	5.90	6.21	3.09	2.07
2001	0.49	1.64	6.51	5.17	9.36	6.32	6.57	4.70	3.40
2002	0.51	2.19	4.56	3.09	8.10	5.91	3.85	3.71	2.76
2003	0.34	2.72	2.85	4.93	6.30	5.02	5.30	3.65	2.63
2004	0.34	2.99	3.99	7.15	8.67	7.08	5.57	4.75	3.46
2005	0.52	1.90	5.05	5.24	7.97	5.78	6.37	4.41	3.13
2006	0.70	2.74	9.75	6.16	6.10	9.97	6.53	5.51	4.14
2007	0.18	2.49	6.07	8.43	14.19	12.76	8.79	7.08	4.73
2008	0.54	2.51	5.75	11.28	11.92	10.34	9.47	7.00	4.83
Whole period	0.42	2.00	4.93	6.09	8.06	7.38	6.29	4.66	3.31
Males									
1999	0.00	0.27	0.28	1.15	2.82	3.90	2.32	1.27	0.82
2000	0.15	0.00	0.29	0.94	2.45	2.45	2.22	1.01	0.68
2001	0.31	0.54	0.58	0.72	1.80	3.15	1.70	1.09	0.82
2002	0.16	0.80	0.87	0.73	2.65	3.23	1.54	1.24	0.91
2003	0.00	1.33	0.58	2.02	3.59	2.72	3.14	1.66	1.16
2004	0.00	0.80	1.74	2.02	2.21	3.39	2.35	1.55	1.11
2005	0.00	0.00	0.85	0.81	3.04	2.23	4.45	1.36	0.81
2006	0.00	0.27	1.13	1.39	3.45	3.86	4.00	1.71	1.09
2007	0.00	0.54	1.11	1.69	6.13	5.85	5.39	2.56	1.60
2008	0.00	0.27	0.55	1.71	3.45	3.86	1.78	1.45	0.97
Whole period	0.06	0.48	0.80	1.32	3.16	3.46	2.89	1.49	1.00

ASR: age specific incidence rate, CR: crude incidence rate, AAR: age adjusted incidence rate (according to Segi's world population).